

**Product Name**

Monoclonal Mouse  
Anti-Blue Fluorescent Protein (BFP) Immunoglobulin,  
clone 7F10

**CAT No.**

MQ 6.104-100

**LOT No.**

TD6.104-11-24-05

**Quantity**

100 µg

Edition: February 1, 2012

**Intended use**

This product is for research use only. NOT for use in diagnostic or therapeutic procedures.

This product is tested for use in enzyme-linked immunosorbent assay (ELISA).

**Reagent provided**

The antibody has been lyophilized in a 10 mM ammonium bicarbonate buffer. Each vial contains 2 mg BSA.

**Isotype**

Mouse IgG1

**Immunogen**

Recombinant EBFP (NCBI accession number AX\_766758 REGION: 1-717, expression vector pGEX-1N), expressed in *E.coli*.

**Specificity**

Specificity has been tested in ELISA (figure 1). Tests for cross reactivity with other fluorescent proteins have not yet been performed. Additional tests for cross reactivity have not yet been performed.

**Purity**

Protein A purified.

**Precautions**

1. For professional users.
2. As with any product derived from biological sources, proper handling procedures should be used.
3. The product may be used in different techniques and in combination with different sample types and materials, therefore each individual laboratory should validate the applied test system.

**Preparation of the antibody**

- Recommended antibody concentration: 0.5 mg/ml (when dissolved at 0.5 mg/ml, the BSA concentration will be 1%).
- Recommended solvent; 100 mM PBS or Tris-HCl, pH 7.0
- Additional sodium azide ( up to 0.05%) is recommended for long term storage.
- For a 0.5 mg/ml antibody concentration in 1% BSA, dissolve in 200 µl buffer.

**NOTE:** Be careful opening the vial since the antibody resides in a vacuum.

**Storage instructions**

Dissolve the antibody and store at 2-8°C.



orders@immunoprecise.com  
www.immunoprecise.com

**Dilution guidelines**

ELISA: 1:(1000 x F) – 1(2000 x F).

**Other applications:** since applications vary, you should determine the optimum working dilution of the product that is appropriate for your specific need.

For the value of the multiplication factor F, see label on vial.

Unless the stability in the actual test system has been established, it is recommended to dilute the product immediately before use.

**Relevance**

Fluorescent proteins, like EBFP, can be used as protein "tags" to study the subcellular localization of proteins and/or their translocation upon stimulation or as markers for transfections in transient and stable expression systems.




Figure 1: Specificity of Anti-Blue Fluorescent Protein (BFP) Immunoglobulin, clone 7F10, determined by ELISA. Antibody fraction (0.5 mg/ml) 900X diluted in PBS containing 0,05% tween-20 and 5% non fat dry milk. Antibody was tested on various recombinant protein substrates i.e. hRpp14-GST (pGEX-2T), hRrp46-HIS10 (pET16b), GST (pGEX-2T) and BSA, 98% (Sigma).