



orders@immunoprecise.com
www.immunoprecise.com

Intended use

This product is for research use only. NOT for use in diagnostic or therapeutic procedures.

This product is tested for use in enzyme-linked immunosorbent assay (ELISA).

Reagent provided

The antibody is supplied in PBS.

Isotype

Human IgG1k

Immunogen

Histidyl-tRNA synthetase. Domain: 1-509 of 509.

Specificity

Specificity has been tested in ELISA (figure 1).

Purity

Protein A purified.

Disclaimer

The antibody is for R&D use only. NOT for use in diagnostic or therapeutic procedures.

Precautions

1. For professional users.
2. As with any product derived from biological sources, proper handling procedures should be used.
3. The product may be used in different techniques and in combination with different sample types and materials, therefore each individual laboratory should validate the applied test system.

Preparation of the antibody

Use antibody as supplied.

Storage/Stability

Store at -20°C. After first time use, store at 4°C. Avoid repeated freeze-thaw cycles.

Application guidelines

ELISA: 1:1000 – 1:5000

Other applications: since applications vary, optimum working dilution of the product should be determined in the appropriate assay.

Unless the stability in the actual test system has been established, it is recommended to dilute the product immediately before use.

Product Name

Monoclonal Human anti-histidyl-tRNA synthetase
Immunoglobulin

CAT No.

MQR 2.3801

LOT No.

19046

Quantity

100 µg

Edition: September 11, 2019

Relevance

Cytoplasmic histidyl-tRNA synthetase, a protein involved in the basic translation machinery taking place in all cells, but also linked to several diseases such as Marie-Charcot disease, Myositis (idiopathic inflammatory myopathies, IIM) and anti-Synthetase Syndrome (aSS). Over evolution, the protein has achieved an additional domain, WHEP, which is believed to carry out non-canonical functions.




Figure 1: Specificity of anti-HARS (MQR2.3801), determined by ELISA. Antibody stock 2.44 mg/ml diluted in PBS containing 0.05% tween-20 and 1% BSA was tested on human Histidyl-tRNA synthetase (in blue) and non-target protein (in red)..

References

- 1) <https://www.uniprot.org/uniprot/P12081>